

General Post-Operative Discharge Instructions

Diet:

Resume your regular diet. Start slow and bland (crackers, toast), then work up to normal foods.
A higher protein intake is recommended to help your body heal. Consider adding protein bars, powder, or shakes to your daily intake.
It is important to drink plenty of fluids so that you do not become dehydrated.

Wound Care:

Leave dressings in place until follow-up
It is okay to remove bandages and shower in 48 hours with indirect water onto your incision (soap and water can lightly run over area, but do not soak area or submerge under water). After your shower, pat area dry and apply new, clean dressings.
Apply antibiotic ointment to the incision 2-3 times daily for one week
Leave any steri-strips (white or brown tape) in place until they fall off on their own. It is okay to trim any edges that start to peel up.
You have dissolvable sutures under your skin and medical skin glue on top. Leave this in place until it falls off on its own. This usually takes 1-3 weeks.
Do NOT smoke or use tobacco products. It WILL impair your healing and increase your complications.

Drain Care (if present):

Empty all drains at least daily, and/or when the bulb gets half full (50mL). Initially you may need to do this several times per day, but it will lessen as your body begins to heal.
Have someone help you with this.
Note how much fluid is in the drain bulb prior to emptying. Record these drain outputs each time (recommend using log sheet provided). Bring recording of daily totals to your follow-up office visit.
To begin, wash your hands with soap and water
Use one hand to hold the tubing tightly near your body (to prevent pulling out the tube). Put a small drop of water or soap on the thumb & index finger of your other hand, pinch the tubing and squeeze the fluid through the tubing toward the bulb.
Open drain bulb cap, record volume, and pour out drainage into a clean measuring cup.
Flush drainage down toilet.
Squeeze the bulb tight and replace the cap to restart suction.
Secure the bulb(s) to clothing, bra, or binder so there is no tugging on the drain tubing.
Anticipate the color to slowly change from a deep red, to a lighter pink/red, and finally a yellowish watery color. It is normal for several large pieces of bloody clot to make their way through the tubing throughout your drain course.
- If drainage becomes heavy (more than 200mL per drain in one day) or has redness or swelling at the incision or drain site(s), notify Dr. Geiger as soon as possible.

Activity:

No strenuous activity or straining until cleared by Dr. Geiger at follow-up.
Do not lift anything heavier than 1-2 pounds (a bottle of water is good estimate. A gallon of milk is too heavy).
Avoid activities that may risk injury to your surgical site.
No driving while taking narcotic pain medications.

Medications: See Patient Visit Summary for list of medications. Do not take pain medications on an empty stomach as this may increase nausea. Avoid Aspirin and NSAIDs (motrin, advil, ibuprofen, aleve) for the first week after surgery, then okay to start taking. Take all of your antibiotic (if prescribed).

Pain medication: You may be prescribed Percocet or Norco for postoperative pain unless allergic or intolerant.

Antibiotic: You may or may not have an antibiotic prescribed

If you are taking narcotic pain medications you may experience some constipation. Drink plenty of water and eat plenty of fiber in your diet. Consider a fiber supplement such as Metamucil®, FiberCon®, or Citrucel®. It is okay to take over the counter stool softeners once or twice daily such as Docusate or Colace® found at your local drug store.

Follow up: Dr. Geiger in 7-10 days. Call office for specific appointment date and time if you do not already have one.

Office contact number: 314-205-6420

Other Instructions:

Should you experience any of these symptoms call the office:

- Fever of 102 degrees Fahrenheit or greater
- Persistent pain, nausea, or vomiting.
- Persistent bleeding.
- Shortness of breath.
- Significant redness or drainage from your incision.
- Significant leg swelling or calf pain
- Any other questions or concerns.

During business hours (8am – 5pm) call:

Chesterfield Plastic Surgery Office: 314-205-6420

On nights or weekends call:

Hospital exchange: 314-364-4176

General Postsurgical Instructions

You may expect to feel dizzy, weak, and drowsy for as long as 24 hours after receiving the medicine that made you sleep (*anesthetic*). The following information pertains to your recovery period for the first 24 hours following surgery.

Do not drive a car, ride a bicycle, participate in physical activities, or take public transportation until you are done taking narcotic pain medicines or as directed by your caregiver.

Do not drink alcohol or take tranquilizers.

Do not take medicine that has not been prescribed by your caregiver.

Do not sign important papers or make important decisions while on narcotic pain medicines. Have a responsible person with you.